

Universal theory of complex SYK models and extremal charged black holes


Subir Sachdev

Chaos and Order: from Strongly Correlated Systems to Black Holes,
Kavli Institute for Theoretical Physics
University of California, Santa Barbara,
December 7, 2018






Wenbo Fu



Yingfei Gu



Grigory Tarnopolsky



1. Quantum matter without quasiparticles:
the complex SYK model
2. Einstein-Maxwell theory of charged
black holes in AdS space
3. Fluctuations, and the Schwarzian
4. Supersymmetric models

1. Quantum matter without quasiparticles:
the complex SYK model

2. Einstein-Maxwell theory of charged
black holes in AdS space

3. Fluctuations, and the Schwarzian

4. Supersymmetric models

The complex SYK model


(See also: the “2-Body Random Ensemble” in nuclear physics; did not obtain the large N limit; T.A. Brody, J. Flores, J.B. French, P.A. Mello, A. Pandey, and S.S.M. Wong, Rev. Mod. Phys. **53**, 385 (1981))

$$H = \frac{1}{(2N)^{3/2}} \sum_{i,j,k,\ell=1}^N U_{ij;k\ell} c_i^\dagger c_j^\dagger c_k c_\ell - \mu \sum_i c_i^\dagger c_i$$

$$c_i c_j + c_j c_i = 0 \quad , \quad c_i c_j^\dagger + c_j^\dagger c_i = \delta_{ij}$$

$$Q = \frac{1}{N} \sum_i c_i^\dagger c_i$$

$U_{ij;k\ell}$ are independent random variables with $\overline{U_{ij;k\ell}} = 0$ and $\overline{|U_{ij;k\ell}|^2} = U^2$
 $N \rightarrow \infty$ yields critical strange metal.




S. Sachdev and J. Ye, PRL **70**, 3339 (1993)

A. Kitaev, unpublished; S. Sachdev, PRX **5**, 041025 (2015)

The complex SYK model

Feynman graph expansion in U_{ijkl} , and graph-by-graph average, yields exact equations in the large N limit:

$$G(i\omega) = \frac{1}{i\omega + \mu - \Sigma(i\omega)} \quad , \quad \Sigma(\tau) = -U^2 G^2(\tau) G(-\tau)$$
$$G(\tau = 0^-) = \mathcal{Q}.$$



The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

- At long times, and at $T = 0$, $G(\tau) \sim |\tau|^{-2\Delta}$ with $\Delta = 1/q$ (\Rightarrow indication there are no quasiparticles)

The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

- At long times, and at $T = 0$, $G(\tau) \sim |\tau|^{-2\Delta}$ with $\Delta = 1/q$ (\Rightarrow indication there are no quasiparticles)
- At general charge Q , there is a spectral symmetry determined by a parameter \mathcal{E} :

$$G(\tau) \sim \begin{cases} -\tau^{-2\Delta} & \tau > 0 \\ e^{-2\pi\mathcal{E}}(-\tau)^{-2\Delta} & \tau < 0 \end{cases}, \quad T = 0$$

The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

- At long times, and at $T = 0$, $G(\tau) \sim |\tau|^{-2\Delta}$ with $\Delta = 1/q$ (\Rightarrow indication there are no quasiparticles)
- At general charge Q , there is a spectral symmetry determined by a parameter \mathcal{E} :

$$G(\tau) \sim \begin{cases} -\tau^{-2\Delta} & \tau > 0 \\ e^{-2\pi\mathcal{E}}(-\tau)^{-2\Delta} & \tau < 0 \end{cases}, \quad T = 0$$

- There is a universal ‘Luttinger relation’ between $-\infty < \mathcal{E} < \infty$ and the total charge $0 < Q < 1$

$$e^{2\pi\mathcal{E}} = \frac{\sin(\pi\Delta + \theta)}{\sin(\pi\Delta - \theta)}$$
$$Q = \frac{1}{2} - \frac{\theta}{\pi} + \left(\Delta - \frac{1}{2}\right) \frac{\sin(2\theta)}{\sin(2\pi\Delta)}$$

A. Georges, O. Parcollet,
and S. Sachdev, PRB **63**,
134406 (2001)
R. Davison, Wenbo Fu,
A. Georges, Yingfei Gu,
K. Jensen, S. Sachdev, PRB
95, 155131 (2017)

The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

- There is a non-vanishing entropy in the zero temperature limit

$$S(T \rightarrow 0) = N s_0 + \dots$$

A. Georges, O. Parcollet, and S. Sachdev, PRB **63**, 134406 (2001)

The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

- There is a non-vanishing entropy in the zero temperature limit

$$S(T \rightarrow 0) = N s_0 + \dots$$

- The saddle point equations imply the relation

$$\frac{ds_0}{dQ} = 2\pi\mathcal{E}$$

Integrating this relation from $s_0 = 0$, $Q = 0$, allows us to compute s_0 as a function of Q .

A. Georges, O. Parcollet, and S. Sachdev, PRB **63**, 134406 (2001)

The complex SYK model

There are 2^N many body levels with energy E . Shown are all values of E for a single cluster of size $N = 12$. The $T \rightarrow 0$ state has an entropy $S_{GPS} = N s_0$, where $s_0 < \ln 2$ is determined by integrating


$$\frac{ds_0}{dQ} = 2\pi\mathcal{E}.$$

At $Q = 1/2$,

$$s_0 = \frac{G}{\pi} + \frac{\ln(2)}{4} = 0.464848\dots$$

where G is Catalan's constant.

GPS: A. Georges, O. Parcollet, and S. Sachdev, PRB **63**, 134406 (2001)



Many-body level spacing $\sim 2^{-N} = e^{-N \ln 2}$

Non-quasiparticle excitations with spacing $\sim e^{-N s_0}$

The complex SYK model

Solution of these equations and corresponding evaluation of the free energy yields the following universal results (*i.e.* all results are *quantitatively* unchanged by adding additional higher q fermion terms):

- At $T > 0$, we obtain a solution with a conformal structure

$$G(\tau) = -A \frac{e^{-2\pi\mathcal{E}T\tau}}{\sqrt{1 + e^{-4\pi\mathcal{E}}}} \left(\frac{T}{\sin(\pi T\tau)} \right)^{1/2}, \quad 0 < \tau < 1/T,$$


where the ‘particle-hole asymmetry’ is determined by \mathcal{E}

A. Georges and O. Parcollet PRB **59, 5341 (1999)**
S. Sachdev, PRX **5, 041025 (2015)**

1. Quantum matter without quasiparticles:
the complex SYK model
2. Einstein-Maxwell theory of charged
black holes in AdS space
3. Fluctuations, and the Schwarzian
4. Supersymmetric models

1. Quantum matter without quasiparticles: the complex SYK model

S. Sachdev, Phys. Rev. Lett. **105**, 151602 (2010)
S. Sachdev, PRX **5**, 041025 (2015)




2. Einstein-Maxwell theory of charged black holes in AdS space

3. Fluctuations, and the Schwarzian

4. Supersymmetric models

1. Quantum matter without quasiparticles: the complex SYK model

S. Sachdev, Phys. Rev. Lett. **105**, 151602 (2010)
S. Sachdev, PRX **5**, 041025 (2015)



2. Einstein-Maxwell theory of charged black holes in AdS space


3. Fluctuations, and the Schwarzian

4. Supersymmetric models

Charged black holes

$$S_{EM} = \frac{1}{2\kappa^2} \int d^{d+2}x \sqrt{-g} \left(\mathcal{R} + \frac{6}{L^2} - \frac{L^2}{g_F^2} F^2 \right)$$

Black hole horizon of radius r_0



Solutions of S_{EM} have metric and gauge field ($F = dA$)

$$ds^2 = -V(r)dt^2 + r^2 d\Omega_d^2 + \frac{dr^2}{V(r)} \quad , \quad A = \mu \left(1 - \frac{r_0^{d-1}}{r^{d-1}} \right) dt$$

$$V(r) = 1 + \frac{r^2}{L^2} + \frac{\Theta^2}{r^{2d-2}} - \frac{M}{r^{d-1}}.$$

where $d\Omega_d^2$ is the metric of the d -sphere. All parameters of the solution are determined in terms of the chemical potential μ , and the Hawking temperature of horizon, T .

Charged black holes

In the $T \rightarrow 0$ limit, at fixed μ , we obtain a charged black hole solution with radius $r_0(T \rightarrow 0, \mu) = R_h$. All properties of this black hole can be expressed in terms of R_h

- The total charge in the black hole is

$$Q = \frac{R_h^{d-1} \sqrt{2d [(d+1)R_h^2 + (d-1)L^2]}}{\kappa^2 g_F L}$$

Charged black holes

In the $T \rightarrow 0$ limit, at fixed μ , we obtain a charged black hole solution with radius $r_0(T \rightarrow 0, \mu) = R_h$. All properties of this black hole can be expressed in terms of R_h

- The total charge in the black hole is

$$Q = \frac{R_h^{d-1} \sqrt{2d [(d+1)R_h^2 + (d-1)L^2]}}{\kappa^2 g_F L}$$

- The Bekenstein-Hawking entropy remains finite as $T \rightarrow 0$ (s_d is the area of the d -dimensional surface of a unit sphere)

$$S(T \rightarrow 0) = s_0 + \dots \quad , \quad s_0 = \frac{2\pi s_d}{\kappa^2} R_h^d$$

Charged black holes

In the $T \rightarrow 0$ limit, at fixed μ , we obtain a charged black hole solution with radius $r_0(T \rightarrow 0, \mu) = R_h$. All properties of this black hole can be expressed in terms of R_h

- In the near-horizon region, we change co-ordinates from r to ζ so that

$$r - R_h = \frac{R_2^2}{\zeta} \quad , \quad R_2 = \frac{LR_h}{\sqrt{d(d+1)R_h^2 + (d-1)^2L^2}}.$$

Then the near-horizon metric becomes $\text{AdS}_2 \times S_d$, with


$$ds^2 = R_2^2 \left[\frac{-dt^2 + d\zeta^2}{\zeta^2} \right] + r_0^2 d\Omega_d^2 \quad , \quad A = \frac{\mathcal{E}}{\zeta} dt.$$

where the dimensionless electric field \mathcal{E} is

$$\mathcal{E} = \frac{g_F LR_h \sqrt{2d [(d+1)R_h^2 + (d-1)L^2]}}{2 [d(d+1)R_h^2 + (d-1)^2L^2]}.$$

Charged black holes

Black hole horizon of radius R_h and entropy s_0



- The entropy s_0 , the charge Q , and the dimensionless electric field \mathcal{E} obey

$$\frac{ds_0}{dQ} = 2\pi\mathcal{E}$$

Charged black holes

In the $T \rightarrow 0$ limit, at fixed μ , we obtain a charged black hole solution with radius $r_0(T \rightarrow 0, \mu) = R_h$. All properties of this black hole can be expressed in terms of R_h

- A probe fermion has a near-horizon Green's function with a conformal structure

$$G(\tau) = -A \frac{e^{-2\pi\mathcal{E}T\tau}}{\sqrt{1 + e^{-4\pi\mathcal{E}}}} \left(\frac{T}{\sin(\pi T\tau)} \right)^{1/2}, \quad 0 < \tau < 1/T,$$

where the ‘particle-hole asymmetry’ is determined by \mathcal{E} . This is identical to the complex SYK model.

1. Quantum matter without quasiparticles:
the complex SYK model

2. Einstein-Maxwell theory of charged
black holes in AdS space

3. Fluctuations, and the Schwarzian

4. Supersymmetric models

Fluctuations

- The fluctuations can be expressed as a path integral over a Green's function $G(\tau_1, \tau_2)$ which is bilocal in time. At frequencies $\ll U$, the action is invariant under reparameterizations $f(\tau)$ and gauge transformations $\phi(\tau)$

$$\tau \rightarrow f(\sigma)$$

$$G(\tau_1, \tau_2) \rightarrow [f'(\sigma_1)f'(\sigma_2)]^{-\Delta} e^{-i(\phi(\sigma_1)-\phi(\sigma_2))} G(\sigma_1, \sigma_2)$$

A. Kitaev, 2015

R. Davison, Wenbo Fu, A. Georges, Yingfei Gu, K. Jensen, S. Sachdev, PRB **95**, 155131 (2017)

Fluctuations

- The saddle-point

$$G(\tau_1 - \tau_2) = -A \frac{e^{-2\pi\mathcal{E}T(\tau_1 - \tau_2)}}{\sqrt{1 + e^{-4\pi\mathcal{E}}}} \left(\frac{T}{\sin(\pi T(\tau_1 - \tau_2))} \right)^{2\Delta}$$

is invariant only under $\text{PSL}(2, \mathbb{R})$ transformations which map the thermal circle onto itself, and an associated gauge transformation

$$\frac{\tan(\pi T f(\tau))}{\pi T} = \frac{a \frac{\tan(\pi T \tau)}{\pi T} + b}{c \frac{\tan(\pi T \tau)}{\pi T} + d}, \quad ad - bc = 1,$$

$$-i\phi(\tau) = 2\pi\mathcal{E}T(\tau - f(\tau))$$

A. Kitaev, 2015

R. Davison, Wenbo Fu, A. Georges, Yingfei Gu, K. Jensen, S. Sachdev, PRB **95**, 155131 (2017)

Fluctuations

Symmetry arguments, and explicit computations, show that the effective action is

$$S_{\text{eff}}[f, \phi] = \frac{NK}{2} \int_0^{1/T} d\tau (\partial_\tau \phi + i(2\pi\mathcal{E}T)\partial_\tau f)^2 - \frac{N\gamma}{4\pi^2} \int_0^{1/T} d\tau \{ \tan(\pi T f(\tau)), \tau \},$$

where $f(\tau)$ is a monotonic map from $[0, 1/T]$ to $[0, 1/T]$, the couplings K , γ , and \mathcal{E} can be related to thermodynamic derivatives and we have used the Schwarzian:

$$\{g, \tau\} \equiv \frac{g'''}{g'} - \frac{3}{2} \left(\frac{g''}{g'} \right)^2.$$

Specifically, an argument constraining the effective at $T = 0$ is

$$S_{\text{eff}} \left[f(\tau) = \frac{a\tau + b}{c\tau + d}, \phi(\tau) = 0 \right] = 0,$$

and this is origin of the Schwarzian.

J. Maldacena and D. Stanford, arXiv:1604.07818;
R. Davison, Wenbo Fu, A. Georges, Yingfei Gu, K. Jensen, S. Sachdev, PRB **95**, 155131 (2017);
A. Gaikwad, L.K. Joshi, G. Mandal, and S.R. Wadia, arXiv:1802.07746

Fluctuations

An *exact* path integral over the effective action leads to the following physical consequences

- The ground state energy with fermion number $NQ + p$ (p integer) varies as

$$E_p = E_0 + \frac{p^2}{2NK}$$

This identifies K with the compressibility $K = dQ/d\mu$ at $T = 0$.

Fluctuations

An *exact* path integral over the effective action leads to the following physical consequences

- The ground state energy with fermion number $NQ + p$ (p integer) varies as

$$E_p = E_0 + \frac{p^2}{2NK}$$

This identifies K with the compressibility $K = dQ/d\mu$ at $T = 0$.

- The low temperature corrections to the entropy at *fixed* Q are

$$S(T \rightarrow 0, Q) = N \left[s_0 + \gamma T + \dots \right] + 2 \ln(U/T) \dots$$

This defines γ as the co-efficient of the linear-in- T specific heat (at fixed Q)

Fluctuations

An *exact* path integral over the effective action leads to the following physical consequences

- The *many*-body density of states, $D(E)$, is related to the grand potential, $\Omega(T)$ by

$$Z = e^{-\Omega(T)/T} = \int_{-\infty}^{\infty} dE D(E) e^{-E/T}$$

We obtain


$$D(E) = \sum_{p=-\infty}^{\infty} e^{2\pi p \mathcal{E}} d(E - E_p)$$

where $N\mathcal{Q} + p$ is the integer fermion number,

$$d(E) \sim \exp(Ns_0) \sinh\left(\sqrt{2N\gamma E}\right), \quad E > 0, \quad e^{-cN} \ll \gamma E \ll N$$

There are exponentially more low energy states than for the quasiparticle case, and $D(E)$ self-averages down to energies exponentially small in N .

The complex SYK model



Many-body level spacing $\sim 2^{-N} = e^{-N \ln 2}$

Non-quasiparticle excitations with spacing $\sim e^{-Ns_0}$

There are 2^N many body levels with energy E . Shown are all values of E for a single cluster of size $N = 12$. The $T \rightarrow 0$ state has an entropy $S_{GPS} = Ns_0$, where $s_0 < \ln 2$ is determined by integrating

$$\frac{ds_0}{dQ} = 2\pi\mathcal{E}.$$

At $Q = 1/2$,

$$s_0 = \frac{G}{\pi} + \frac{\ln(2)}{4} = 0.464848\dots$$

where G is Catalan's constant.

GPS: A. Georges, O. Parcollet, and S. Sachdev, PRB **63**, 134406 (2001)

Fluctuations

An *exact* path integral over the effective action leads to the following physical consequences

- The *many*-body density of states, $D(E)$, is related to the grand potential, $\Omega(T)$ by

$$Z = e^{-\Omega(T)/T} = \int_{-\infty}^{\infty} dE D(E) e^{-E/T}$$

We obtain

$$D(E) = \sum_{p=-\infty}^{\infty} e^{2\pi p \mathcal{E}} d(E - E_p)$$

where $N\mathcal{Q} + p$ is the integer fermion number,

$$d(E) \sim \exp(Ns_0) \sinh\left(\sqrt{2N\gamma E}\right), \quad E > 0, \quad e^{-cN} \ll \gamma E \ll N$$

There are exponentially more low energy states than for the quasiparticle case, and $D(E)$ self-averages down to energies exponentially small in N .

Fluctuations


An *exact* path integral over the effective action leads to the following physical consequences

- At charge $NQ + p$, the prefactor of the $\sinh(\sqrt{2N\gamma(E - E_p)})$ term is

$$\exp[Ns_0(Q) + 2\pi p\mathcal{E}] \approx \exp[Ns_0(Q + p/N)]$$

using

$$\frac{ds_0}{dQ} = 2\pi\mathcal{E}$$



The Schwarzian theory and black holes

- Reparameterization invariance is a defining property of Einstein's theory of gravity
- In imaginary time, AdS_2 is the homogeneous hyperbolic space: two-dimensional surface of constant negative curvature. Its metric is invariant under $SL(2, \mathbb{R})$

$ds^2 = (d\tau^2 + d\zeta^2)/\zeta^2$ is invariant under

$$\tau' + i\zeta' = \frac{a(\tau + i\zeta) + b}{c(\tau + i\zeta) + d} \text{ with } ad - bc = 1.$$



The Schwarzian theory and black holes

- Reparameterization invariance is a defining property of Einstein's theory of gravity
- In imaginary time, AdS_2 is the homogeneous hyperbolic space: two-dimensional surface of constant negative curvature. Its metric is invariant under $SL(2, \mathbb{R})$

Semiclassical fluctuations about the saddle-point of Einstein-Maxwell theory of a charged black holes in $d \geq 2$ spatial dimensions lead to the same Schwarzian+phase theory of fluctuations.



P. Nayak, A. Shukla, R.M. Soni, S.P. Trivedi, and V. Vishal, arXiv:1802.09547

U. Moitra, S. P. Trivedi, and V. Vishal, arXiv:1808.08239

P. Chaturvedi, Yingfei Gu, Wei Song, Boyang Yu, arXiv:1808.08062

A. Gaikwad, L.K. Joshi, G. Mandal, and S.R. Wadia, arXiv:1802.07746

The Schwarzian theory and black holes

- The Einstein-Maxwell theory leads to the following parameters for the Schwarzian+phase theory

$$K = \left. \frac{dQ}{d\mu} \right|_{T=0} = \frac{2(d-1)s_d R_h^{d-3} [d(d+1)R_h^2 + (d-1)^2 L^2]}{(d+1)g_F \kappa^2}$$

$$S(T \rightarrow 0, Q) = s_0 + \gamma T + \dots$$

$$\gamma = \frac{4\pi s_d L^2 R_h^{d+1}}{\kappa^2 (d(d+1)R_h^2 + (d-1)^2 L^2)}.$$



1. Quantum matter without quasiparticles:
the complex SYK model

2. Einstein-Maxwell theory of charged
black holes in AdS space

3. Fluctuations, and the Schwarzian

4. Supersymmetric models

SYK with $\mathcal{N} = 2$ SUSY

$$Q = i \sum_{1 \leq i < j < k \leq N} C_{ijk} c_i c_j c_k$$

$$H = QQ^\dagger + Q^\dagger Q = \sum_{i,j,k,\ell}^N U_{ij;kl} c_i^\dagger c_j^\dagger c_k c_\ell$$

C_{ijk} are independent random variables with $\overline{C_{ijk}} = 0$
and $\overline{|C_{ijk}|^2} = 2U/N^2$.

SYK with $\mathcal{N} = 2$ SUSY

$$Q = i \sum_{1 \leq i < j < k \leq N} C_{ijk} c_i c_j c_k$$

$$H = QQ^\dagger + Q^\dagger Q = \sum_{i,j,k,\ell}^N U_{ij;kl} c_i^\dagger c_j^\dagger c_k c_\ell$$

C_{ijk} are independent random variables with $\overline{C_{ijk}} = 0$
and $\overline{|C_{ijk}|^2} = 2U/N^2$.

There is a $U(1)_R$ charge

$$Q_R = \frac{1}{3} \sum_i \left(c_i^\dagger c_i - \frac{1}{2} \right)$$

The smallest absolute values of are $Q_R = 0$ (for N even) and $Q_R = \pm 1/6$ (for N odd) and we focus on these cases: $\mathcal{N} = 2$ SUSY is preserved by the ground states. Cases with extensive Q_R charge, $\lim_{N \rightarrow \infty} Q_R/N \neq 0$, break SUSY and require a more complex analysis.

SYK with $\mathcal{N} = 2$ SUSY

- There is an *exact*, exponentially large degeneracy of zero energy states for any C_{ijk} . Consequently, $e^{Ns_0} \equiv g(N, \mathcal{Q}_R)$ is an integer for all N (there was no such requirement for the non-SUSY case). The exact non-zero values for small \mathcal{Q}_R are

$$\begin{aligned} g(N, 0) &= 2 \times 3^{N/2-1} \quad , \quad g(N, \pm 1/3) = 3^{N/2-1} \quad , \quad \text{for } N \text{ even} \\ g(N, \pm 1/6) &= 3^{(N-1)/2} \quad , \quad \text{for } N = 3 \text{ mod } 4 \\ g(N, \pm 1/6) &= 3^{N/2-1} \quad , \quad g(N, \pm 1/2) = 1 \text{ or } 3 \quad , \quad \text{for } N = 1 \text{ mod } 4 \end{aligned}$$


There are no zero energy states for other values of \mathcal{Q}_R . So in the large N limit, the entropy $s_0 = (1/2) \ln 3$ for $\lim_{N \rightarrow \infty} \mathcal{Q}_R/N = 0$.

SYK with $\mathcal{N} = 2$ SUSY


- Fluctuations are described by a super-Schwarzian theory for $\lim_{N \rightarrow \infty} Q_R/N = 0$. The density of states has delta function at zero energy, and a continuum contribution at non-zero energies, both of order $e^{s_0 N}$. There is a gap between the ground states and the excited states $\sim 1/N$ for even N .

D. Stanford and E. Witten, arXiv:1703.04612

$\mathcal{N} = 2$ (N even)





$\mathcal{N} = 2$ (N odd)



SYK with $\mathcal{N} = 2$ SUSY

- Fluctuations are described by a super-Schwarzian theory for $\lim_{N \rightarrow \infty} Q_R/N = 0$. The density of states has delta function at zero energy, and a continuum contribution at non-zero energies, both of order $e^{s_0 N}$. There is a gap between the ground states and the excited states $\sim 1/N$ for even N .

D. Stanford and E. Witten, arXiv:1703.04612



SYK with $\mathcal{N} = 2$ SUSY

- Fluctuations are described by a super-Schwarzian theory for $\lim_{N \rightarrow \infty} Q_R/N = 0$. The density of states has delta function at zero energy, and a continuum contribution at non-zero energies, both of order $e^{s_0 N}$. There is a gap between the ground states and the excited states $\sim 1/N$ for even N .

D. Stanford and E. Witten, arXiv:1703.04612

- Similar results are expected to $\mathcal{N} = 2$ black holes with AdS_2 horizons obtained from string theory (Sen, Dabholkar, Gomes, Murthy ...).

SYK with $\mathcal{N} = 2$ SUSY

- Fluctuations are described by a super-Schwarzian theory for $\lim_{N \rightarrow \infty} \mathcal{Q}_R/N = 0$. The density of states has delta function at zero energy, and a continuum contribution at non-zero energies, both of order $e^{s_0 N}$. There is a gap between the ground states and the excited states $\sim 1/N$ for even N .

D. Stanford and E. Witten, arXiv:1703.04612

- Similar results are expected to $\mathcal{N} = 2$ black holes with AdS_2 horizons obtained from string theory (Sen, Dabholkar, Gomes, Murthy ...).
- For $\lim_{N \rightarrow \infty} \mathcal{Q}_R/N \neq 0$, we expect that the density of states of the generic complex SYK model will apply.

Quantum matter without quasiparticles

- Planckian dynamics is realized in the ‘solvable’ SYK models
- Black holes thermalize in a time $\sim \hbar/(k_B T_H)$, where T_H is the Hawking temperature.
- A Schwarzian theory of a time reparameterization mode, with $SL(2, \mathbb{R})$ symmetry, describes the quantum dynamics of
 - the SYK models
 - black holes with near-extremal AdS_2 horizons