


# Decoupling and the MSSM


## Higgs mass

LHC—the first part of the journey  
KITP Conference

Howard E. Haber  
July 8, 2013



The Kavli Institute for  
Theoretical Physics  
University of California, Santa Barbara



## Outline

- The Higgs sector of the MSSM
- The neutral MSSM Higgs masses
  - tree level results
  - including the radiative corrections
- Impact of the discovery of a Higgs boson of mass 125 GeV
  - Constraints on MSSM mass parameters
  - Implications for specific SUSY-breaking models
- Questioning the robustness of the MSSM Higgs mass prediction
- The one-loop MSSM Higgs mass revisited
  - a consistent low energy definition of  $\tan \beta$
  - the decoupling of high mass scale physics
- The Higgs mass prediction in the seesaw-extended MSSM
  - decoupling of the heavy right-handed neutrino sector
  - effective field theory estimates of the Higgs mass shift
- Conclusions

## The Higgs sector of the MSSM

The Higgs sector of the MSSM is a two Higgs doublet model, whose scalar potential and Yukawa couplings are constrained by supersymmetry (SUSY). The scalar potential of the MSSM is:

$$V = (m_d^2 + |\mu|^2) H_d^{i*} H_d^i + (m_u^2 + |\mu|^2) H_u^{i*} H_u^i - m_{ud}^2 (\epsilon^{ij} H_d^i H_u^j + \text{h.c.}) \\ + \frac{1}{8} (g^2 + g'^2) [H_d^{i*} H_d^i - H_u^{j*} H_u^j]^2 + \frac{1}{2} g^2 |H_d^{i*} H_u^i|^2,$$

where  $\mu$  is a supersymmetric Higgsino mass parameter and  $m_d^2$ ,  $m_u^2$ ,  $m_{ud}^2$  are soft-SUSY-breaking masses.

Minimizing the Higgs potential, the neutral components of the Higgs fields acquire vacuum expectation values (vevs),  $\langle H_d^0 \rangle = v_d/\sqrt{2}$  and  $\langle H_u^0 \rangle = v_u/\sqrt{2}$ , where  $v^2 \equiv v_d^2 + v_u^2 = 4m_W^2/g^2 = (246 \text{ GeV})^2$ . The ratio of the two vevs is an important parameter of the model:

$$\tan \beta \equiv \frac{v_u}{v_d}.$$

## Tree-level neutral MSSM Higgs masses

The CP-even Higgs bosons  $h$  and  $H$  are eigenstates of the squared-mass matrix

$$\mathcal{M}_0^2 = \begin{pmatrix} m_A^2 \sin^2 \beta + m_Z^2 \cos^2 \beta & -(m_A^2 + m_Z^2) \sin \beta \cos \beta \\ -(m_A^2 + m_Z^2) \sin \beta \cos \beta & m_A^2 \cos^2 \beta + m_Z^2 \sin^2 \beta \end{pmatrix}.$$

The eigenvalues of  $\mathcal{M}_0^2$  are the squared-masses of the two CP-even Higgs scalars

$$m_{H,h}^2 = \frac{1}{2} \left( m_A^2 + m_Z^2 \pm \sqrt{(m_A^2 + m_Z^2)^2 - 4m_Z^2 m_A^2 \cos^2 2\beta} \right),$$

and  $\alpha$  is the angle that diagonalizes the CP-even Higgs squared-mass matrix. It follows that

$$m_h \leq m_Z |\cos 2\beta| \leq m_Z.$$

If this tree-level mass inequality were more generally satisfied, then the MSSM would be ruled out today!

## The radiatively-corrected mass of $h^0$


The tree-level inequality,  $m_h \leq m_Z$ , is significantly modified by quantum corrections. The mass of the lightest CP-even Higgs boson of the MSSM can be shifted due to an incomplete cancellation from loops of particles and their superpartners [H.E. Haber and R. Hempfling (1991); Y. Okada, M. Yamaguchi and T. Yanagida (1991); J.R. Ellis, G. Ridolfi and F. Zwirner (1991)]:




$$m_h^2 \lesssim m_Z^2 + \frac{3g^2 m_t^4}{8\pi^2 m_W^2} \left[ \ln \left( \frac{M_S^2}{m_t^2} \right) + \frac{X_t^2}{M_S^2} \left( 1 - \frac{X_t^2}{12M_S^2} \right) \right],$$

where  $X_t \equiv A_t - \mu \cot \beta$  governs stop mixing and  $M_S^2$  is the geometric mean of the squared-masses of the top-squarks  $\tilde{t}_1$  and  $\tilde{t}_2$  (which are the mass-eigenstates derived from the interaction eigenstates,  $\tilde{t}_L$  and  $\tilde{t}_R$ ). Here, only the leading one-loop log and leading squark mixing contributions are exhibited.



The state-of-the-art computation includes the full 1-loop result, all the significant 2-loop contributions, some of the leading 3-loop terms, and renormalization-group improvements.



## Implications of the observed Higgs state with $m_h \simeq 125$ GeV




Contour plot of  $m_h$  in the  $M_S$  vs.  $X_t$  plane, with  $\tan \beta = 30$  and  $M_Q = M_U = M_S$ . The solid curve is  $m_h = 125$  GeV with  $m_t = 173.2$  GeV. The band around the solid curve corresponds to  $m_h = 125 \pm 2$  GeV. The dashed lines correspond to varying  $m_t$  from 172–174 GeV. Taken from P. Draper, P. Meade, M. Reece and D. Shih, Phys. Rev. D **85**, 095007 (2012).




Contour plot of  $X_t$  in the plane of physical stop masses  $(m_{\tilde{t}_1}; m_{\tilde{t}_2})$ . Here  $X_t$  is fixed to be the minimum positive (left) or negative (right) solution to  $m_h = 125$  GeV. Taken from P. Draper, P. Meade, M. Reece and D. Shih (2012).

Even without assumptions about the SUSY-breaking mechanism, the observed Higgs mass tends to push some MSSM parameters into the multi-TeV regime. This provides significant tension with naturalness constraints. The tension is exacerbated in specific SUSY breaking models.



CMSSM in the focus point region with dark matter constraints  
from J.L. Feng, K.T. Matchev and D. Sanford (2012)



Gauge-Mediated SUSY breaking with  $A_t = 0$  at the high scale  
from P. Draper, P. Meade, M. Reece and D. Shih (2012)

But, is the MSSM Higgs mass prediction reliable? Could it potentially be modified by new physics that lies significantly above the TeV scale?

The Higgs mass prediction relies on decoupling—very heavy states that do not receive their masses from electroweak symmetry breaking should have a negligible impact on the Higgs mass prediction.

In 2011, S. Heinemeyer, M.J. Herrero, S. Penaranda and A.M. Rodriguez-Sanchez [JHEP **1105**, 063 (2011)] analyzed corrections to the MSSM Higgs mass in the seesaw-extended MSSM. Their results seemed to suggest that contributions from the right-handed neutrino sector could alter the Higgs mass prediction by a few GeV. If true, one might accommodate the observed Higgs mass more comfortably within some SUSY-breaking scenarios.

Patrick Draper and I argue in arXiv:1304.6103 [hep-ph] that this interpretation is not correct. Decoupling of heavy-scale physics does hold as expected, and the impact of the right-handed neutrino sector of the seesaw-extended MSSM is utterly negligible and thus can be safely ignored in the Higgs mass prediction.

## The MSSM Higgs mass at one-loop

Although rarely displayed, the complete 1-loop expressions for the pole masses of the CP-even neutral MSSM Higgs bosons are given by:

$$m_h^2 = (m_h^2)^{\text{tree}} - \frac{\sqrt{2}A_h}{v} s_{\beta-\alpha} - \Sigma_{ZZ}(m_Z^2) s_{\beta+\alpha}^2 + \Sigma_{hh}(m_h^2) \\ - \Sigma_{AA}(m_A^2) c_{\beta-\alpha}^2 + \Sigma_{GG}(0) s_{\beta-\alpha}^2 - 4m_Z^2 c_\beta^2 s_{\beta+\alpha} c_{\beta+\alpha} \delta \tan \beta ,$$

$$m_H^2 = (m_H^2)^{\text{tree}} - \frac{\sqrt{2}A_H}{v} c_{\beta-\alpha} - \Sigma_{ZZ}(m_Z^2) c_{\beta+\alpha}^2 + \Sigma_{HH}(m_H^2) \\ - \Sigma_{AA}(m_A^2) s_{\beta-\alpha}^2 + \Sigma_{GG}(0) c_{\beta-\alpha}^2 + 4m_Z^2 c_\beta^2 s_{\beta+\alpha} c_{\beta+\alpha} \delta \tan \beta ,$$

where  $s_{\beta-\alpha} \equiv \sin(\beta - \alpha)$ ,  $c_{\beta-\alpha} \equiv \cos(\beta - \alpha)$ , etc., and the Higgs mixing angle  $\alpha$  is determined implicitly via its tree-level relation,

$$\frac{m_A^2}{m_Z^2} = -\frac{c_{\beta+\alpha} s_{\beta+\alpha}}{c_{\beta-\alpha} s_{\beta-\alpha}} .$$

For consistency of the one-loop approximation, the arguments of the self-energies are evaluated by their *tree-level* values.

The  $\Sigma$  functions for the scalars (vectors) are the real parts of one loop self-energies (proportional to  $g^{\mu\nu}$ ). Here, the on-shell scheme is used in defining the renormalized physical boson masses.

In terms of the one-loop tadpoles,  $A_u$  and  $A_d$ , of the hypercharge  $\pm 1$  neutral Higgs fields (which are determined by the requirement that they cancel the corresponding tree-level tadpoles), we have defined,

$$A_h \equiv A_u \cos \alpha - A_d \sin \alpha, \quad A_H \equiv A_u \sin \alpha + A_d \cos \alpha.$$

These are related to the Goldstone self-energy,

$$\sqrt{2}v\Sigma_{GG}(0) = A_H c_{\beta-\alpha} + A_h s_{\beta-\alpha},$$

which follows from the requirement that the one-loop Goldstone boson mass vanishes.

In order to make use of the above formulae, we must decide on a method for fixing the  $\tan \beta$  counterterm, denoted above by  $\delta \tan \beta$ .

## A low energy definition of $\tan \beta$

Consider the seesaw-extended MSSM. How does the heavy right-handed neutrino sector affect the predicted values of  $m_h^2$  and  $m_H^2$ ? The answer depends on the definition of  $\tan \beta$ . If you define  $\tan \beta$  based on physical quantities that can be measured by experiments that probe the TeV scale, then the effects of the heavy right-handed neutrino sector are completely negligible.

Here is a simple example of a low energy definition of  $\tan \beta$ . We call this scheme the Higgs mass (HM) scheme. In this scheme, we use  $m_H$  as an input parameter in place of  $\tan \beta$ . In this case,

$$m_h^2 = m_A^2 + m_Z^2 - m_H^2 + A_h(m_h^2) + A_H(m_H^2) - \Sigma_{ZZ}(m_Z^2) - \Sigma_{AA}(m_A^2) - \Sigma_{GG}(0) ,$$

a result originally obtained by M. Berger in 1990. [Theoretical issues associated with the definition of  $\tan \beta$  have also been considered by A. Freitas and D. Stockinger (2002).]

In the HM scheme, the  $\tan \beta$  counterterm is obtained by setting  $m_H^2 = (m_H^2)^{\text{tree}}$  in the one-loop expression for  $m_H^2$ , which *defines*  $\tan \beta$  in terms of the physical parameters  $m_Z$ ,  $m_H$  and  $m_A$ ,

$$(\delta \tan \beta)_{\text{HM}} = \frac{1}{4m_Z^2 c_\beta^2 s_{\beta+\alpha} c_{\beta+\alpha}} \left( \frac{\sqrt{2}A_H}{v} c_{\beta-\alpha} + \Sigma_{ZZ}(m_Z) c_{\beta+\alpha}^2 - \Sigma_{HH}(m_H) \right. \\ \left. + \Sigma_{AA}(m_A) s_{\beta-\alpha}^2 - \Sigma_{GG}(0) c_{\beta-\alpha}^2 \right).$$

A second possible scheme is to define  $\tan \beta$  via the decay  $A^0 \rightarrow \tau\tau$ . The  $\tan \beta$  counterterm would then depend on the measured partial width.

An alternative strategy: define  $\tan \beta$  via the  $\overline{\text{DR}}$  scheme. Not surprisingly, the effects of the high-scale physics do not decouple. In this case,  $\tan \beta_{\overline{\text{DR}}}$  is not directly a physical parameter. If the non-decoupling physics is known, then one can evaluate both the radiatively-corrected Higgs mass and a second low-energy observable in terms of  $\tan \beta_{\overline{\text{DR}}}$ . If  $\tan \beta_{\overline{\text{DR}}}$  is then eliminated, then one prediction remains and all effects of high-scale physics decouple.

Can  $\overline{\text{DR}}$  be modified to respect decoupling? The  $\text{m}\overline{\text{DR}}$  scheme of Heinemeyer et al. attempts to do this by removing by hand terms that are logarithmically sensitive to the high energy scale. But, this procedure fails to remove constant terms induced by the high-scale physics that can be of order a few GeV. Such terms are absent in the physical schemes described previously.

We have developed an alternative scheme that automatically removes both the large logarithms and the constant terms induced by the high-scale physics. After renormalizing the vevs,

$$v_u \rightarrow \mathcal{Z}_{H_u}^{-1/2} v_u = v_u \left(1 + \frac{1}{2} \delta \mathcal{Z}_{H_u}\right), \quad v_d \rightarrow \mathcal{Z}_{H_d}^{-1/2} v_d = v_d \left(1 + \frac{1}{2} \delta \mathcal{Z}_{H_d}\right),$$

the  $\tan \beta$  counterterm, defined by  $\tan \beta \rightarrow \tan \beta - \delta \tan \beta$  is given by:

$$\delta \tan \beta \equiv \frac{1}{2} (\delta \mathcal{Z}_{H_d} - \delta \mathcal{Z}_{H_u}) \tan \beta .$$

We now introduce the decoupling scheme (DEC) to fix the wave function renormalization,

$$(\delta \mathcal{Z}_{H_d})_{\text{DEC}} = \left. \frac{d\Sigma_{HH}(p^2)}{dp^2} \right|_{\alpha=0, p^2=0}, \quad (\delta \mathcal{Z}_{H_u})_{\text{DEC}} = \left. \frac{d\Sigma_{hh}(p^2)}{dp^2} \right|_{\alpha=0, p^2=0} .$$

The large logarithms and the constant terms induced by the high-scale physics are manifestly removed. Although  $\tan \beta_{\text{DEC}}$  is not directly a physical parameter, its definition is completely insensitive to the high-scale physics.

## The seesaw extended MSSM

Introduce a right-handed neutrino superfield  $N$  and a superpotential

$$W = \mu H_d H_u + y_\nu L H_u N - y_l L H_d R + \frac{1}{2} m_M N N .$$

Add soft SUSY-breaking terms,

$$V_{\text{soft}} = m_{\tilde{R}}^2 \tilde{N}^* \tilde{N} + (y_\nu A_\nu H_U^0 \tilde{\nu}_L \tilde{N}^* + m_M B_\nu \tilde{N} \tilde{N} + \text{h.c.}) .$$

As a result, one obtains the seesaw neutrino mass matrix,

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & m_D \\ m_D & m_M \end{pmatrix} ,$$

where  $m_D \equiv y_\nu v_u$ . The CP-even/odd (+/−) sneutrino mass matrices are given by:

$$\mathcal{M}_{\tilde{\nu}_\pm}^2 = \begin{pmatrix} m_{\tilde{L}}^2 + m_D^2 + \frac{1}{2} m_Z^2 \cos 2\beta & m_D (A_\nu - \mu \cot \beta \pm m_M) \\ m_D (A_\nu - \mu \cot \beta \pm m_M) & m_{\tilde{R}}^2 + m_D^2 + m_M^2 \pm 2B_\nu m_M \end{pmatrix} ,$$

where  $m_{\tilde{L}}^2$  is the usual soft-breaking mass for the left-handed sneutrinos present in the MSSM.

## Decoupling of the right-handed neutrino sector in the one-loop expression for $m_h$

For simplicity, set  $\mu = A_\nu = B_\nu = 0$  and fix  $m_{\tilde{L}} = m_{\tilde{R}} \equiv m_S$ . Then expand to first order in  $m^2/m_M^2$ , where  $m \in \{m_Z, m_S, m_D\}$ , and to leading order in powers of  $m_Z$  (the latter is equivalent to setting  $\alpha = \beta - \frac{1}{2}\pi$ ). At leading-log order, the squared-mass of the lightest CP-even Higgs boson is shifted relative to its tree level value in the HM and DEC schemes by an amount:

$$\left(\Delta m_h^2\right)_{\text{HM}} \simeq \frac{g^2}{48\pi^2 c_W^2} m_Z^2 \log \frac{m_S}{m_Z} - \frac{g^2 m_D^4 m_S^2}{4\pi^2 c_W^2 m_M^2 m_Z^2 \sin^2 \beta} \log \frac{m_M}{m_S},$$

$$\left(\Delta m_h^2\right)_{\text{DEC}} \simeq \frac{g^2}{48\pi^2 c_W^2} m_Z^2 \cos^2 2\beta \log \frac{m_S}{m_Z} - \frac{g^2 m_D^4 m_S^2}{4\pi^2 c_W^2 m_M^2 m_Z^2} \log \frac{m_M}{m_S},$$

where  $c_W \equiv \cos \theta_W = m_W/m_Z$ . The second term in both expressions above, which is generated by the right-handed neutrino sector, yields a correction  $\Delta m_h \sim m_\nu^2/m_h$  (where  $m_\nu \sim m_D^2/m_M$ ) and is thus utterly negligible.

The difference in the two results can be accounted for by the different definitions of  $\tan \beta$ . Indeed,

$$\tan \beta_{\text{HM}} = \tan \beta_{\text{DEC}} + \delta \tan \beta_{\text{HM}} - \delta \tan \beta_{\text{DEC}},$$

implies that

$$\left( \Delta m_h^2 \right)_{\text{DEC}} - \left( \Delta m_h^2 \right)_{\text{HM}} \simeq -2m_Z^2 \cos^2 \beta \sin 4\beta \left[ \delta \tan \beta_{\text{HM}} - \delta \tan \beta_{\text{DEC}} \right].$$

If we evaluate  $\delta \tan \beta$  in the two schemes employing the same approximations used to obtain  $\Delta m_h^2$  above, we obtain:

$$\frac{\delta \tan \beta_{\text{HM}} - \delta \tan \beta_{\text{DEC}}}{\tan \beta} \simeq c_{2\beta} \left( \frac{g^2}{96\pi^2 c_W^2} \log \frac{m_S}{m_Z} - \frac{g^2 m_D^4 m_S^2}{32\pi^2 c_W^2 m_M^2 m_Z^4 s_\beta^4} \log \frac{m_M}{m_S} \right).$$

Substituting this expression above then reproduces the difference in the two expressions for  $\Delta m_h^2$  previously obtained.

We now compare the mass shift in the DEC scheme to that of the  $\overline{\text{DR}}$  scheme for defining the  $\tan\beta$  counterterm (the leading results in the HM scheme are identical). The potential non-decoupling terms that arise in the Higgs mass computation, which are of  $\mathcal{O}(m_D^2)$ , cancel precisely in the DEC scheme, where

$$\delta \tan\beta_{\overline{\text{DEC}}} = \frac{g^2 m_D^2}{32\pi^2 c_W^2 m_Z^2 \sin 2\beta} \left( \frac{1}{\epsilon} - \gamma + \log 4\pi - \log \frac{m_M^2}{Q^2} + 1 \right),$$

where  $Q$  is the arbitrary mass scale of dimensional regularization. In the  $\overline{\text{DR}}$  scheme, only  $\epsilon^{-1} - \gamma + \log 4\pi$  is retained in  $\delta \tan\beta$ . Hence,

$$\left( \Delta m_h^2 \right)_{\overline{\text{DR}}} \simeq \left( \Delta m_h^2 \right)_{\text{DEC}} + \frac{g^2 m_D^2}{8\pi^2 c_W^2} \cos^2 \beta \cos 2\beta \left( \log \frac{m_M^2}{Q^2} - 1 \right),$$

Even if one sets  $Q = m_M$ , which removes the large logs by hand, one is left with a term of  $\mathcal{O}(m_D^2)$  can yield a Higgs mass shift as large as a few GeV. This is a remnant of the right-handed neutrino scale and must also be removed to restore the decoupling behavior.

## Effective field theory estimates of the Higgs mass shift


Suppose that we integrate out the right-handed neutrino and sneutrino at the right-handed neutrino mass threshold. Above this scale, the running of the Higgs quartic coupling is supersymmetric, but the TeV-scale soft mass splits the scalar and fermion states, leading to a logarithmic correction to the quartic coupling  $\lambda$  from the right-handed sneutrino bubble diagram:

$$\Delta m_h^2 = 2v^2 \Delta\lambda \sim \frac{m_D^4}{v^2} \log \frac{m_{\tilde{N}}^2}{m_N^2} \sim \frac{m_D^4 m_S^2}{v^2 m_M^2} .$$

This term is  $m_M$ -suppressed and has no log enhancement. In addition to direct contributions to the Higgs quartic coupling, we also generate an approximately supersymmetric dimensional-five coupling,

$$\Delta W \sim \frac{y_\nu^2}{m_M} LHLH .$$

The dimension-five coupling affects the running of the quartic coupling and the running of  $\tan \beta$  when supersymmetry is broken via the diagrams:



The dominant contribution comes from the sneutrino diagram,

$$\frac{\partial \lambda}{\partial \log Q^2} \approx \frac{y_\nu^4 m_S^2 \sin^4 \beta}{8\pi^2 m_M^2}.$$

Running the quartic coupling from  $m_M$  to  $m_S$ , we obtain at leading log order,

$$\Delta m_h^2 = -\frac{m_D^4 m_S^2}{2\pi^2 v^2 m_M^2} \log \frac{m_M}{m_S},$$

matching the result previously obtained in the DEC scheme. The running of  $\tan \beta$  in the HM scheme introduces an additional contribution to  $\Delta m_h^2$ .

## Large SUSY-breaking in the right-handed neutrino sector?

One might be tempted to consider the possibility of choosing large values for the SUSY-breaking parameters  $m_{\tilde{R}}^2$  and  $B_\nu$ . An effective field theory estimate shows that

$$\Delta m_h^2 \sim \frac{m_D^4}{v^2} \log \left( \frac{m_M^2 + m_{\tilde{R}}^2}{m_M^2} \right),$$

and

$$\Delta m_h^2 \sim \frac{m_D^4}{v^2} \log \left( \frac{m_M^2 - B_\nu^2}{m_M^2} \right),$$

respectively, which would yield a Higgs mass shift of order  $\Delta m_h \sim m_D^4 / (v^2 m_h)$ .

However, naturalness constraints suggest that  $m_{\tilde{R}}$  should not be larger than other soft-SUSY-breaking parameters, and  $B_\nu$  cannot be larger than about  $10^3 m_{\tilde{\nu}_L}$  in order to avoid generating too large a one-loop mass for neutrinos via  $\tilde{\nu}_L$ - $\tilde{\nu}_R$  mixing [Y. Grossman and H.E. Haber (1997)].

## Conclusions

- It is possible to accommodate  $m_h \sim 125$  GeV within the MSSM. However, this value strongly suggests that the relevant SUSY-breaking parameters must be at least of  $\mathcal{O}(1 \text{ TeV})$  or higher, which provides tension with expectations of naturalness. This tension is often exacerbated in specific SUSY-breaking models.
- The MSSM predictions for the masses of the neutral CP-even Higgs bosons are robust. Potential contributions to these masses due to additional physics at a very high mass scale are strongly suppressed (decoupling!).
- As an example, the contribution in the seesaw-extended MSSM from the heavy right-handed neutrino sector to the one-loop corrected MSSM Higgs masses (when expressed in terms parameters that can be determined by TeV-scale observables) is utterly negligible.